

# 2008 Area Source Emissions Inventory Methodology 699 – OTHER (MISCELLANEOUS PROCESSES)

## I. Purpose

This document describes the Area Source Methodology used to estimate emissions of carbon monoxide (CO), carbon dioxide (CO<sub>2</sub>), nitrogen oxides (NO<sub>x</sub>), fine particulate matter less than 10 microns (PM<sub>10</sub>), volatile organic compounds (VOC), and sulfur oxides (SO<sub>x</sub>) from other processes not accounted for in other source categories in the San Joaquin Valley Air Basin. An area source is a collection of similar emission units within a geographic area (ie., a County). Area sources collectively represent individual sources that are small and numerous, and that may not have inventoried as specific point, mobile, or biogenic sources. The California Air Resources Board (CARB) has grouped these individual sources with other like sources into area source categories. These source categories are grouped in such a way that they can be estimated collectively using one methodology.

# II. Applicability

This Area Source Methodology applies to facilities that are identified by the following Category of Emission Source (CES) code and Reconciliation Emission Inventory Code (REIC):

Table 1. Emission inventory code

CES	REIC	Description
90480	699-995-0000-0000	All Other Unspecified Sources

#### **III.** Point Source Reconciliation

Emissions from the area source inventory and point source inventory are reconciled against each other to prevent double counting. This is done using relationships created by the California Air Resources Board (CARB) between the area source REIC and the point sources' Standard Industry Classification (SIC) code and emissions process Source Category Code (SCC) combinations. Currently, this source category does not reconcile to any processes within our point source inventory.

Rev. Date: 19 January 2010 Rev. By: J. Hart

# IV. Methodology Description

The purpose of this source category is to estimate emissions from processes not accounted for in other source categories. At this time, the District has not identified any other sources of emissions that are not accounted for elsewhere. Therefore, the area source emissions for this category will be set to zero.

# V. Activity Data

Not applicable.

#### **VI.** Emission Factors

Not applicable.

#### VII. Emissions Calculations

#### A. Assumptions

Not applicable.

## B. Sample Calculations

Not applicable.

# VIII. Temporal Variation

Not applicable.

# IX. Spatial Variation

Not applicable.

## X. Growth Factor

Not applicable.

## XI. Control Level

Not applicable.

Rev. Date: 19 January 2010

Rev. By: J. Hart

## XII. CARB Chemical Speciation

CARB has developed organic gas profiles in order to calculate reactive organic gasses (ROG), volatile organic compounds (VOC) or total organic gas (TOG) given any one of the three values. For each speciation profile, the fraction of TOG that is ROG and VOC is given. The organic gas profile codes can also be used to lookup associated toxics. CARB's organic gas speciation profile for other process sources is presented in the table below:

Table 2. CARB organic gas speciation profile for other process sources.

	CARB	Fractions		
Profile Description	Organic Gas Profile#	ROG	voc	
All Other Unspecified Sources	600	0.6986	0.6986	

CARB has developed particulate matter speciation profiles in order to calculate particulate matter (PM), particulate matter with a diameter less than or equal to 10 microns (PM $_{10}$ ) or particulate matter with a diameter less than or equal to 2.5 microns (PM $_{2.5}$ ) given any one of the three values. For each speciation profile, the fraction of PM that is PM $_{10}$  and PM $_{2.5}$  is given. The particulate matter profile codes can also be used to lookup associated toxics. CARB's particulate matter speciation profile for other process sources is presented in the table below:

Table 3. CARB particulate matter speciation profile for other process sources.

Profile Description	CARB PM	Fractions		
Profile Description	Profile#	PM <sub>10</sub>	PM <sub>2.5</sub>	
All Other Unspecified Sources	900	0.7	0.42	

Rev. Date: 19 January 2010

Rev. By: J. Hart

# XIII. Assessment Of Methodology

This source category is not currently used by the District.

### XIV. Emissions

Following is the 2008 area source emissions inventory for REIC 699-995-0000-0000 estimated by this methodology. Emissions are reported for each county in the District.

Table 4. Area source emissions for REIC 699-995-0000-0000 (2008).

County	Emissions (tons/year)					
County	NOx	CO	SOx	VOC <sup>(1)</sup>	PM <sub>10</sub>	PM <sub>2.5</sub> <sup>(2)</sup>
Fresno	0.00	0.00	0.00	0.00	0.00	N/A
Kern	0.00	0.00	0.00	0.00	0.00	N/A
Kings	0.00	0.00	0.00	0.00	0.00	N/A
Madera	0.00	0.00	0.00	0.00	0.00	N/A
Merced	0.00	0.00	0.00	0.00	0.00	N/A
San Joaquin	0.00	0.00	0.00	0.00	0.00	N/A
Stanislaus	0.00	0.00	0.00	0.00	0.00	N/A
Tulare	0.00	0.00	0.00	0.00	0.00	N/A
TOTAL	0.00	0.00	0.00	0.00	0.00	N/A

<sup>(1)</sup> The District only reports ROG to CARB. As noted in Section XII, ROG is the same as VOC.

Following is the 2008 point source emissions inventory for REIC 699-995-0000-0000 as reported to the District by our permit holders. Emissions are reported for each county in the District.

Table 5. Point source emissions for REIC 699-995-0000-0000 (2008).

County	Emissions (tons/year)					
County	NOx	CO	SOx	VOC <sup>(1)</sup>	PM <sub>10</sub>	PM <sub>2.5</sub> <sup>(2)</sup>
Fresno	0.00	0.00	0.00	0.00	0.00	N/A
Kern	0.00	0.00	0.00	0.00	0.00	N/A
Kings	0.00	0.00	0.00	0.00	0.00	N/A
Madera	0.00	0.00	0.00	0.00	0.00	N/A
Merced	0.00	0.00	0.00	0.00	0.00	N/A
San Joaquin	0.00	0.00	0.00	0.00	0.00	N/A
Stanislaus	0.00	0.00	0.00	0.00	0.00	N/A
Tulare	0.00	0.00	0.00	0.00	0.00	N/A
TOTAL	0.00	0.00	0.00	0.00	0.00	N/A

<sup>(1)</sup> The District only reports ROG to CARB. As noted in Section XII, ROG is the same as VOC.

Rev. Date: 19 January 2010

Rev. By: J. Hart

<sup>(2)</sup> At this time, the District does not calculate PM<sub>2.5</sub> emissions. PM<sub>2.5</sub> emissions can be estimated using the speciation profiles found in Section XII.

<sup>(2)</sup> At this time, the District does not calculate PM<sub>2.5</sub> emissions. PM<sub>2.5</sub> emissions can be estimated using the speciation profiles found in Section XII.

Following is the 2008 total unreconciled (point source plus area source) emissions inventory for REIC 699-995-0000-0000. Emissions are reported for each county in the District.

Table 6. Total emissions for REIC 699-995-0000-0000 (2008).

County	Emissions (tons/year)					
County	NOx	CO	SOx	VOC <sup>(1)</sup>	PM <sub>10</sub>	PM <sub>2.5</sub> <sup>(2)</sup>
Fresno	0.00	0.00	0.00	0.00	0.00	N/A
Kern	0.00	0.00	0.00	0.00	0.00	N/A
Kings	0.00	0.00	0.00	0.00	0.00	N/A
Madera	0.00	0.00	0.00	0.00	0.00	N/A
Merced	0.00	0.00	0.00	0.00	0.00	N/A
San Joaquin	0.00	0.00	0.00	0.00	0.00	N/A
Stanislaus	0.00	0.00	0.00	0.00	0.00	N/A
Tulare	0.00	0.00	0.00	0.00	0.00	N/A
TOTAL	0.00	0.00	0.00	0.00	0.00	N/A

<sup>(1)</sup> The District only reports ROG to CARB. As noted in Section XII, ROG is the same as VOC.

Following is the net change in total unreconciled emissions between this update (2008 inventory year) and the previous emissions year (2007 inventory year) for REIC 699-995-0000-0000. The change in emissions are reported for each county in the District.

Table 7. Net emissions change for REIC 699-995-0000-0000 (2007-2008).

County	Emissions (tons/year)						
County	NOx	CO	SOx	VOC <sup>(1)</sup>	PM <sub>10</sub>	PM <sub>2.5</sub> <sup>(2)</sup>	
Fresno	0.00	0.00	0.00	0.00	0.00	N/A	
Kern	0.00	0.00	0.00	0.00	-8.15	N/A	
Kings	0.00	0.00	0.00	0.00	0.00	N/A	
Madera	0.00	0.00	0.00	0.00	0.00	N/A	
Merced	0.00	0.00	0.00	0.00	0.00	N/A	
San Joaquin	0.00	0.00	0.00	0.00	0.00	N/A	
Stanislaus	0.00	0.00	0.00	0.00	0.00	N/A	
Tulare	0.00	0.00	0.00	0.00	0.00	N/A	
TOTAL	0.00	0.00	0.00	0.00	0.00	N/A	
(4) TI D'							

<sup>(1)</sup> The District only reports ROG to CARB. As noted in Section XII, ROG is the same as VOC.

Rev. Date: 19 January 2010 Rev. By: J. Hart

<sup>(2)</sup> At this time, the District does not calculate  $PM_{2.5}$  emissions.  $PM_{2.5}$  emissions can be estimated using the speciation profiles found in Section XII.

<sup>(2)</sup> At this time, the District does not calculate PM2.5 emissions. PM2.5 emissions can be estimated using the speciation profiles found in Section XII.

# XV. Revision History

2010. This is a new District methodology.

# XVI. Update Schedule

In an effort to provide inventory information to CARB and other District programs and maximize limited resources, the District has developed an update cycle based on emissions within the source category as shown in the table below:

Table 8. Area source update frequency criteria.

Total Emissions (tons/day)	Update Cycle (years)		
<=1	4		
>1 and <= 2.5	3		
>2.5 and <=5	2		
>5	1		

Since no emissions sources have been identified for *All Other Unspecified Sources*, this source category will be reviewed for activity every four years.

## XVII. References

Not applicable.

Rev. Date: 19 January 2010

Rev. By: J. Hart